

SPACE ENVIRONMENT TECHNOLOGIES

Space Weather Division

SDO-EVE workshop

USU CASS ionospheric research
SET solar ops requirements JB2008
USU/SWC GAIM, and SpaceWx app

SDO EVE Workshop


October 7, 2010

W. Kent Tobiska

Space Environment Technologies

Utah State University Space Weather Center

<http://SpaceWx.com>



Topics

Research needs

Operational needs

Distributed networks as basis for operational systems

JB2008

GAIM

SpaceWx app

SET forecasts



SPACE ENVIRONMENT TECHNOLOGIES


Space Weather Division

SDO-EVE workshop

Research needs





S2K-EVE comparison





S2K-EVE comparison ratio






TDIM: USU Time Dependent Ionospheric Model

Principle Inputs controlled by $F_{10.7}$, IMF, K_p (or A_p):

- Convection electric field [Heppner and Maynard, 1987]
- Auroral oval [Hardy et al., 1987]
- Neutral atmosphere - MSIS [Hedin, 1987]
- Thermospheric Wind - HWM [Hedin et al., 1991]

Outputs: Profiles of NO^+ , O_2^+ , O^+ from 100 to 800 km





Previously, the TDIM has always used the same default solar EUV spectrum for calculating the production rates in sunlight, based on a 37-bin scheme [Hinteregger, 1977]. Now we are investigating the use of spectra that are high-resolution both in time steps and in wavelength bins.



TDIM: Log Ne 2010/125 comparison



SC21REFW spectrum

Log Ne day 125 TDIM old default spectrum





S2K spectrum

Log Ne day 125 Kent's single 39-bin spectrum



EVE spectrum

Log Ne day 125 SDO-EVE time-dependent spectra






TDIM: E- and F-region using 3 solar inputs

SC21REFW spectrum

S2K spectrum


EVE spectrum






TDIM: TEC comparison using 3 solar inputs


SC21REFW spectrum




S2K spectrum




EVE spectrum






Research Recommendations

1. able to incorporate EVE measurements and model spectra into physics-based ionosphere models (TDIM) to produce reasonable results; lessons from TDIM can be extended to IFM
2. differences between the time resolved EVE data and daily S2K model provide a way to improve the representation of solar energy relevant to the ionosphere for periods when no data are available; need to recalibrate S2K to EVE include 1 minute time series data representation in S2K
3. use GAIM ionosphere as baseline for specific dates, events, and time series; its a path for understanding how to improve the solar and ionosphere models; need to archive GAIM runs to capture events



Research Recommendations

4. need calibrated daily solar EVE data as a data product to allow validation of global parameters in models such as heating efficiencies and to allow validation of solar EUV/FUV heating and photoionization rates in TDIM
5. during flare events, time series calibrated EVE data can provide inputs to TDIM for GAIM comparison for photoionization rate validation, for solar zenith angle validation, and for unit optical depth validation of species transport due to dynamics
6. solar minimum to solar maximum daily EVE data used in TDIM/GAIM comparison can help understand climate-scale problems related to thermospheric cooling; can help understand mechanisms that may have accounted for extremely low ionosphere in solar minimum



Research Recommendations

7. TDIM has high ionosphere densities during low solar activity and this needs study
8. need to begin pulling in data more consistently and regularly from LASP
9. need for a spectrum plot – perhaps use small dots to represent EVE data points throughout the day to get the range of variation, do the mean value of the EVE data for the day, and that will provide the ability to compare with S2K and old spectrum for the day




SPACE ENVIRONMENT TECHNOLOGIES


Space Weather Division

SDO-EVE workshop

Operational needs




Distributed networks as basis for operational systems




Distributed Network versus Clustered System

- ◆ The basis for contemporary operational needs comes from the formation and use of distributed networks
- ◆ A distributed network differs from the more common rack-mount, clustered operational system
- ◆ In both cases a Tier 1 central server/database exist
 - Distributed networks use Tier 1 to link with servers/algorithms at geographically dispersed locations
 - Rack-mount clusters use Tier 1 to operate on local algorithms




What is a Distributed Network?

- ◆ Distributed networks link automated systems of models across dispersed geographical locations
- ◆ Outputs from one automated system are linked to the inputs of another automated system to form an integrated, but geographically dispersed, operational network
- ◆ Distributed networks allow developers to maintain versioning and proprietary control over their models while exchanging data via automated systems
- ◆ Distributed networks use dynamic metadata which allows traceability for all I/O requests from models, customers, or users



What is Dynamic Metadata?

- ◆ User devices or systems access data products by making discrete, secure, formatted requests to a server using application software
- ◆ Devices or systems receive back formatted data for use by their own algorithms
- ◆ Example – a user algorithm may want altitude-dependent electron densities for a particular time, latitude, longitude; request is sent to database and the extracted/returned data objects are provided “just-in time”



Smart phones are the tip of the iceberg for distributed networks





- ◆ automated systems provide immediate data to mobile smart phones; they connect satellite and ground-based data streams with algorithms to quickly process the measurements into geophysical data, incorporate those data into operational space physics models, and generate images, plots, and alerts that can be viewed on smart phones
- ◆ institutional teams now collaborate in prototyping and operational development processes; SpWx community is going through transition from research models to operations by proofing products in real-time via smart phones
- ◆ SDO/EVE data often has a few minutes latency between GEO measurement and SpaceWx app display





Distributed network example

solar-ionosphere
coupling affects the
quality of HF
communications





SPACE ENVIRONMENT TECHNOLOGIES

Space Weather Division


SDO-EVE workshop

JB2008




JB2008 overview

JB2008 output




neutral
atmosphere

JB2008 solar inputs




solar EUV
& FUV
indices

JB2008 geo- mag inputs




a_p and Dst
indices




Uses of JB2008 in CIRA08, ISO 14222, and HASDM ASW

- **CIRA08** is the **scientific publication** describing the Earth's neutral atmosphere, especially above 120 km, and is compiled by authors in the COSPAR CIRA task group
- **ISO 14222** is the Earth atmosphere density >120 km **international standard** developed by ISO TC20/SC14/WG4 project leads
- **HASDM ASW** is the **operational system** used to specify the JB2008 neutral atmosphere densities for debris avoidance and space situational awareness by Air Force Space Command. Solar and geomagnetic indices are provided with 72-hour forecast in real-time system.



EUV heating and photoionization





Compare fluxes






Table A. Solar indices related to atmospheric heating


Index	IS 21348 Spectral category	IS 21348 Spectral sub-category	Wavelength range (nm)	Solar source temperature region ¹	Solar source feature ¹	Atmosphere absorption (unit optical depth, km) ²	Terrestrial atmosphere absorption (thermal region) ²
*F _{10.7}	Radio	Radio	10.7E7	Transition region, cool corona	Active region	90-500	Thermosphere with 1-day lag; 9.8% daily variability contribution
*S _{10.7}	UV	EUV	26-34	Chromosphere, corona	Active region, plage, network	200-300	Thermosphere with 1-day lag; 74.1% daily variability contribution
*M _{10.7}	UV	FUV	160	Photosphere-lower chromosphere	SRC	95-110	Lower thermosphere with 2-day lag; 10.3% daily variability contribution
MgII _{cwr}	UV	MUV	280 ³	Chromosphere	Active region, plage, network	200-300	Thermosphere
*Y _{10.7}	X-rays and UV	X-rays+H Lyman-α	0.1-0.8, 121	Chromosphere, transition region, hot corona	Active region, plage, network	85-100	Mesopause-lower thermosphere with 5-day lag; 5.8% daily variability contribution
H Lyα	UV	H Lyman-α	121	Transition region, chromosphere	Active region, plage, network	85-100	Mesopause-lower thermosphere
X _{b10}	X-rays	X-rays	0.1-0.8	Hot corona	Active region background	85-100	Mesopause-lower thermosphere

*Index or proxy is used in the JB2008 model exospheric temperature equation.

¹Vernazza *et al.*, 1976; Vernazza *et al.*, 1981.

²Banks, P. and G. Kockarts, 1973.

³The *h* & *k* lines at the band center are chromospheric and are referenced to blackbody continuum wings at edges of bandpass.



Indices' characteristics

- By expressing the proxies and indices in common units, their contribution to the daily density variability in the JB2008 exospheric temperature equation for T_c can be determined; the $F_{10.7}$ contribution to T_c daily variability is 9.8%, $S_{10.7}$ is 74.1%, $M_{10.7}$ is 10.3%, and $Y_{10.7}$ is 5.8%


Table B. Characteristics of daily JB2008 solar indices

Index or proxy	Observing facility	Instrument	Observation time frame	Measurement cadence	Measurement latency	Operational availability
$F_{10.7}$	Penticton ground observatory	Radio telescope	1947-2009	3 times/day	Up to 24 hours	yes
$S_{10.7}$	SOHO, GOES	SEM, EUVS	1996-2009	15-second	Up to 24 hours	(a)
$M_{10.7}$	NOAA-16,17,18, SORCE, ERS-2	SBUV, SOLSTICE, GOME	1991-2009	2 times/day	Up to 24 hours	yes
$Y_{10.7}$	GOES-12, UARS, SORCE, TIMED	XRS, SOLSTICE (2), SEE	1991-2009	1-minute, 16 times/day	Up to 10 minutes, up to 48 hours	(b)

(a) SOHO/SEM is a NASA research instrument but provides daily irradiances on an operational cadence; GOES 13 EUVS B channel makes measurements in the same bandpass as SOHO SEM.

(b) GOES XRS is a NOAA operational instrument whereas TIMED/SEE and SORCE/SOLSTICE are NASA research instruments providing daily irradiances on an operational measurement cadence.

(c) UARS/SOLSTICE stopped in 2005; SORCE/SOLSTICE intends to provide data for several years.




Indices for different solar activity conditions

Table A4. Reference values for intermediate- and short-term solar variability


	Case 1: Intermediate-term (81 days)			Case 2: Short-term (27 days high activity)			Case 3: Short-term (27 days low activity)		
Daily	Low	Moderate	High	Low	Moderate	High	Low	Moderate	High
$F_{10.7}$	65	120	225	90	165	280	80	105	145
$S_{10.7}$	60	120	215	105	135	185	85	100	120
$M_{10.7}$	60	115	215	95	135	185	80	95	115
$Y_{10.7}$	50	115	180	110	150	185	90	110	135

Daily updated values are found at the JB2008 menu link on the SET <http://spacewx.com> website



F10 proxy

- First daily measurements began February 14, 1947 – **high value for legacy time series**
- Measured by DRAO National Research Council Solar Radio Monitoring Programme in Penticton
- Observations of flux density values made at 17, 20 and 23 UTC each day
- 20 UTC observed values (not 1 AU) archived at World Data Center were used in JB2008 derivation
- Physical units are $\times 10^{-22} \text{ W m}^{-2} \text{ Hz}^{-1}$ and their numerical value is used without the multiplier and called solar flux units (sfu); **this has become defacto standard unit for solar indices**
- $F_{10.7}$ a useful proxy for the combination of **chromospheric, transition region, and coronal** solar EUV emissions modulated by bright solar active regions whose energies at Earth are deposited in the thermosphere
- It originates mostly in the solar cool (low) corona by electrons in thermal free-free (bremsstrahlung) emission in the vicinity of sunspots and in widely distributed areas associated with the hot complexes of solar activity




F10 proxy


- The dependence on few processes, combined with its localized formation in the cool corona, i.e., a region that is closely coupled with magnetic structures responsible for creating the XUV–EUV irradiances, makes this a **good generalized solar proxy** for thermospheric heating
- The running 81-day centered smoothed $F_{10.7}$ values, using the moving boxcar method, are referred to as F_{81}
- The observed archival daily $F_{10.7}$ values and their 81-day running center-smoothed values, F_{81} , with a **1-day lag** are used in JB2008
- The 1-day lag had the best correlation with satellite-derived density residuals
- A linear regression is used with daily $F_{10.7}$ to scale and **report all other JB2008 solar indices in units of sfu**
- $F_{10.7}$ is the **recognized historical EUV proxy** and, by reporting other proxies or indices in sfu, it is very easy to qualitatively identify similarities and differences between them
- For solar energy inputs, it is **desirable to have solar indices and proxies that vary differently through time**; this strategy of using multiple solar indices has significantly improved the accuracy of density modeling in JB2008



F10 proxy




$F_{10.7}$ daily and
81-day
smoothed
values for use
by the JB2008
model from
January 1,
1997 to
January 1,
2009



S10 index

- $S_{10.7}$ index is an activity indicator of the integrated 26–34 nm solar irradiance measured by the Solar Extreme-ultraviolet Monitor (SEM) instrument on the NASA/ESA Solar and Heliospheric Observatory (SOHO) satellite
- SEM has been making observations since December 16, 1995 for the 26–34 nm solar EUV emission with 15-second time resolution in this first order broadband wavelength range
- $S_{10.7}$ index is created by first normalizing the data, then converting it to sfu via a first degree polynomial fit with $F_{10.7}$; spikes from abnormal flares and missing data were excluded from the fitting vectors; normalization is achieved using a mean value = 1.9955×10^{10} photons $\text{cm}^{-2} \text{s}^{-1}$ for the time frame December 16, 1995 to June 12, 2005 (solar cycle 23)
- Corrections to $S_{10.7}$ are made and identified with versions: versions 3.0–3.9 are used by JB2006 and after June 12, 2005 include a slight long-term trend removal to ensure that similar values at the minima of solar cycles 22 and 23 were achieved
- For JB2008, ($S_{10.7}$ v4.0 and higher) a new derivation was completed using the equation as follows:
$$S_{10.7} = (-2.90193) + (118.512) * (\text{SOHO_SEM}_{26_34} / 1.9955 \times 10^{10})$$




S10 index


- Chromospheric He II at 30.4 nm and coronal Fe XV at 28.4 nm dominate the broadband SEM 26-34 nm irradiances but that bandpass includes contributions from other chromospheric, transition region, and coronal lines
- When the SOHO SEM and TIMED SEE 26–34 nm integrated data are compared, there are differences in the time series particularly during active solar conditions; it is possible that the SOHO SEM measurements are slightly contaminated with 2nd order emissions from the coronal 17.1 nm Fe IX line that have not been removed; however, this topic needs further investigation
- The energy in this bandpass principally comes from solar active regions, plage, and network
- Once the photons reach the Earth, they are deposited (absorbed) in the terrestrial thermosphere mostly by atomic oxygen above 200 km
- We use the daily $S_{10.7}$ index and its 81-day running center-smoothed values, S_{81} , with a 1-day lag (the best correlation with satellite density residuals) in JB2008
- We infer the 1-day lag is consistent with the average atomic oxygen thermal conduction timescale in the thermosphere above 180 km



S10 index




$S_{10.7}$ (v4.0)
daily and 81-
day smoothed
values for use
by the JB2008
model from
January 1,
1997 to
January 1,
2009




M10 proxy

- The $M_{10.7}$ index is derived from the Mg II core-to-wing ratio that originated from the NOAA series operational satellites, e.g., NOAA-16,-17,-18, which host the Solar Backscatter Ultraviolet (SBUV) spectrometer
- Although the NOAA data are from operational satellites, the SORCE/SOLSTICE and ERS-2/GOME research satellites also make the Mg II cwr measurements
- The 280 nm solar spectral band contains photospheric continuum and chromospheric line emissions; the Mg II h and k lines at 279.56 and 280.27 nm are chromospheric in origin while the weakly varying wings or continuum longward and shortward of the core line emission are photospheric in origin; the instruments from all satellites observe both features
- The ratio of the Mg II variable core lines to the nearly non-varying wings is calculated to provide a measure of chromospheric solar active region emission that is theoretically independent of instrument sensitivity change through time; however, long-term changes can occur in the index if instrument wavelength calibrations change in-flight or the solar incidence angle into the instrument changes



M10 proxy

- The daily Mg II core-to-wing ratio (cwr) has historically been provided through the NOAA Space Weather Prediction Center (SWPC); SET has developed and provides an operational Mg II cwr data product ($\text{MgII}_{\text{cwr_SET}}$) available at the Products menu link of <http://spacewx.com> that uses the NOAA-16,-17-18, SORCE/SOLSTICE, and ERS-2/GOME data sources
- SET uses the DeLand algorithm to create the index
- The Mg II cwr is an especially good proxy for some solar FUV and EUV emissions; it well represents photospheric and lower chromospheric solar FUV Schumann-Runge Continuum emission near 160 nm that maps into lower thermosphere heating due to O_2 photodissociation; since a 160 nm solar FUV emission photosphere index is not produced operationally, the $\text{MgII}_{\text{cwr_SET}}$ proxy is used and modified into the $M_{10.7}$ index for comparison with the other solar indices
- This derivation is performed by finding the relationship between long-term (multiple solar cycle) daily $\text{MgII}_{\text{cwr_SET}}$ and $F_{10.7}$ using a first-degree polynomial fit to produce a coefficient set that can translate Mg II cwr into sfu; the result is $M^*_{10.7}$




M10 proxy


- Next, a correction is added for the decline of solar cycle 23 to account for NOAA 16 instrument degradation that may be related to its diffuser screen illumination geometry changing with time; this cause is unconfirmed; the correction is accomplished by using another first degree polynomial fit between a trend ratio and day number starting 2448542.0 JD (October 12, 1991 12:00 UT) near the peak of solar cycle 22; the trend ratio is formed from the 365-day center smoothed $M^*_{10.7}$ divided by the 365-day center smoothed $F_{10.7}$
- The v4.0 formulation of the $M_{10.7}$ index in sfu is: $M_{10.7} = [-2107.6186 + (8203.0537) * (MgII_{cwr_SET})] + [(M^*_{10.7}) * (1.2890589 + (-8.3777235 \times 10^{-5}) * x - 1)]$
- The day number $x = 0, 1, 2, \dots$ with $x = 0$ equivalent to starting on 2448542.0 JD
- The daily $M_{10.7}$ and its 81-day running center-smoothed values, M_{81} , are used with a 2-day lag in JB2008 as a proxy for the Schumann-Runge continuum FUV emission
- A 2-day lag time is appropriate for $M_{10.7}$, which represents O_2 photodissociation, recombination, conduction, and transport processes at the 95–110 km level
- We infer this lag is consistent with the average molecular oxygen dissociation and thermal conduction timescale in the thermosphere above 95 km, although eddy and turbulent conduction processes may play a role



M10 proxy




$M_{10.7}$ (v4.0)
daily and 81-day smoothed
values for use
by the JB2008
model from
January 1,
1997 to
January 1,
2009




Y10 index

- The $XL_{10.7}$ index was developed as a candidate index for the JB2006 model but was unused; while developing the JB2008 model, it was determined that a thermospheric energy contribution to satellite drag from the 80–95 km region was significantly correlated with the composite $XL_{10.7}$ solar index
- Solar X-rays in the 0.1–0.8 nm wavelength range come from the cool and hot corona and are typically a combination of both very bright solar active region background that varies slowly (days to months) plus flares that vary rapidly (minutes to hours), respectively
- The photons arriving at Earth are absorbed in the lower thermosphere to mesopause and (85–100 km) by molecular oxygen (O_2) and molecular nitrogen (N_2) where they ionize those neutral constituents to create the ionospheric D-region
- The X-ray Spectrometer (XRS) instrument is part of the instrument package on the GOES series operational spacecraft. The GOES/XRS provides the historical through current epoch 0.1–0.8 nm solar X-ray emission data with a 1-minute cadence and as low as 5-minute latency. These data, which are particularly useful for flare detection, are continuously reported by NOAA SWPC at their website of <http://www.swpc.noaa.gov/>



Y10 index

- SET uses the GOES/XRS 0.1–0.8 nm data for an index of the solar X-ray active region background, without the flare component, for operational use; this is called the X_{b10} index and is used to represent the daily energy that is deposited into the mesosphere and lower thermosphere
- While the 0.1-0.8 nm X-rays are a major energy source in these atmospheric regions during high solar activity, they relinquish their dominance to another emission that reaches the same optical depth, i.e., the competing hydrogen (H) Lyman- α emission that is the major energy source in this atmosphere region during moderate and low solar activity
- Lyman- α is created in the solar upper chromosphere and transition region and demarcates the EUV from the FUV spectral regions; it is formed primarily in solar active regions, plage, and network; the photons, arriving at Earth, are absorbed in the mesosphere and lower thermosphere where they dissociate nitric oxide (NO) and participate in water (H₂O) chemistry; Lyman- α is regularly observed by the SOLSTICE instrument on the UARS and SORCE satellites as well as by the SEE instrument on TIMED and the EVE instrument on SDO




Y10 index

- Since these two solar emissions are competing drivers to the mesosphere and lower thermosphere, we have developed a composite solar index of the X_{b10} and Lyman- α ; it does not contain a flare component and is weighted to represent mostly X_{b10} during solar maximum and to represent mostly Lyman- α during moderate and low solar activity
- A normalized F_{81} , F_{81norm} , consisting of the 81-day centered smoothed $F_{10.7}$ divided by its mean value for the time frame of January 1, 1991 through February 16, 2008 is used as the weighting function and multiplied with the X_{b10} and Lyman- α (Lya) expressed as ratios to their solar maximum values; the resulting index is called $Y_{10.7}$ in sfu
- $Y_{10.7} = F_{81norm} * X10 + [(1 - F_{81norm}) * L10]$
- $L10 = -88.3926 + (3.35891 \times 10^{-10} * Lya) + (2.40481 \times 10^{-22} * Lya^2)$
- $X10 = [(-42.5991 + (0.533669 * X_{b10}))]$
- This daily index has a 5-day lag strongly correlated with the satellite drag density residuals after modeled density variations due to the other solar indices were removed; the 81-day running center-smoothed values, Y_{81} , are also used with the 5-day lag
- We infer the 5-day lag is consistent with the average molecular oxygen and molecular nitrogen thermal conduction timescales in the lower thermosphere above 85 km, although eddy and turbulent conduction may also play a role







Y10 index




$Y_{10.7}$ (v4.0)
daily and 81-day
smoothed
values for use
by the
JB2008
model from
January 1,
1997 to
January 1,
2009



Comparison between indices




$F_{10.7}$, $S_{10.7}$,
 $M_{10.7}$, $Y_{10.7}$
(v4.0) monthly
minimum,
mean, and
maximum
values for use
by the JB2008
model from
January 1,
1997 to
January 1,
2009





a_p index

- The a_p index reports the amplitude of planetary geomagnetic activity for a given day (Mayaud, 1980); the official a_p values are calculated at the GeoForschungsZentrum Potsdam Adolf-Schmidt-Observatory for Geomagnetism in Germany
- It is translated from the K_p index, which is derived from geomagnetic field measurements made at several locations around the world
- The daily A_p is obtained by averaging the eight 3-hour values of a_p for each day. The U.S. Air Force Weather Agency also calculates an estimated a_p from a different, smaller set of stations than are used in calculating the official a_p values
- The AFWA a_p index values are available through several products issued by NOAA SPWC
- Daily A_p and 3-hour a_p indices were used in early orbit analyses and it was determined that the time scales of geomagnetically-induced variability represented by these indices were on the order of a few hours




Ap index





Dst index

- The Disturbance Storm Time (Dst) index is an indicator of the strength of the storm-time ring current in the inner magnetosphere
- During the main phase of geomagnetic storms, the ring current becomes highly energized and produces a southward-directed magnetic field perturbation at low latitudes on the Earth's surface; this is opposite to the normal northward-directed main field
- The quick-look Dst index is calculated hourly and released through the World Data Center (WDC) in Kyoto, Japan using measurements from four off-equatorial magnetic observatories
- Dst is an 'absolute' index and is reported in units of nT; magnetic observatory data are required for its calculation; magnetic observatories are specially designed and carefully operated facilities that provide stable-baseline magnetometer data over long periods of time; typically, an observatory supports the operation of fluxgate, proton-precession, and declination-inclination (theodolite) magnetometer systems; measurements from the various sensor systems can be combined to produce data that are extremely accurate
- Traditionally, four stations have been used for the Dst calculation: Hermanus (HER), South Africa; Kakioka (KAK), Japan; Honolulu (HON), Hawaii; and San Juan (SJG), Puerto Rico






Dst index

- Most magnetic storms begin with sharp decreases (southward-directed negative values) in Dst, called the storm sudden commencement, in response to increased solar wind pressure. Following a southward turning of the interplanetary magnetic field, Dst decreases as ring current energy increases during the storm's main phase. During the recovery phase the ring current energy decreases and Dst increases until the storm's end when the magnetic field perturbation has ceased
- Use of Dst as an index for the energy deposited in the thermosphere during magnetic storms is more accurate than the use of the a_p index. This is because Dst has higher temporal resolution with an ability to segregate storm phases with their corresponding magnitudes. On the other hand, the 3-hour a_p index is an indicator of general magnetic activity over the Earth and responds primarily to currents flowing in the ionosphere and only secondarily to magnetospheric variations. The a_p index is derived from measurements by observatories at high latitudes that can be blind to energy input during large storms and it can underestimate the effects of storms on the thermosphere
- An algorithm for determining the storm events was developed in JB2008 that locates the temporal start, minimum, recovery slope change, and final end of the storm as reflected in the Dst index




Dst index







Geomagnetic comparisons




The a_p , Dst, and dTc geomagnetic, ring current, and delta temperature indices for use by the JB2008 model in solar cycle 23




The a_p , Dst, and dTc geomagnetic, ring current, and delta temperature indices for use by the JB2008 model in a storm period between November 19-22, 2003




JB2008 output



JB2008 density at 400 km altitude for low solar and quiet geomagnetic activity on July 14, 2010



JB2008 mean air density with altitude for low, moderate, and high long- and short-term solar and geomagnetic activity



JB2008 Operations Recommendations

JB2008: solar irradiance surrogates for redundancy stream (atmospheric drag)

- **S10 index** (26-34 nm EUV calibrated to SOHO SEM dataset) (F2-region ionosphere, >180 km thermosphere)
- **M10 proxy** (160 nm AlA irradiance or Mg II index) (E-region ionosphere, 100-120 km thermosphere)
- **Y10 index** (XRS_L index is the 0.1-0.8 nm modeled X-ray; Lyman- α) (D-region ionosphere, 80-90 km mesosphere)
- **F10 proxy** (E10 index of 1-105 nm integrated flux) (residual ionosphere and thermosphere)
- time granularity (1-hour), cadence (1-hour)




SPACE ENVIRONMENT TECHNOLOGIES

Space Weather Division

SDO-EVE workshop



GAIM



GAIM submodels require solar inputs

Ionosphere Forecast Model (IFM) provides the E-, F-region physics

- Drivers to IFM are empirical models
 - Neutral densities (MSIS 1986 uses F10.7 or E10.7)
 - Thermospheric winds (HWM 1990 uses F10.7 or E10.7)
 - Electric fields convection (HM 1987)
 - Particle precipitation (Hardy 1987)
- Torr & Torr 37+2=39 wavelength groups of solar spectral irradiances used for photoionization of neutral species





ABBYNormal provides D-region physics

- **GOES XRS B (0.1-0.8 nm)** used for photoionization of neutral species

GAIM output is used for HF communication and GPS uncertainty derivative products




GAIM output







GAIM derivatives

HF propagation time series





HF global propagation for 24 cities at 15 MHz



Point-to-point ray-trace

GPS uncertainty maps





GAIM Operations Recommendations

GAIM: solar irradiance surrogates and spectral formats (HF communication and GPS uncertainty)

- **E10 index** (driver to IFM submodels)
- **39 wavelength group** format (driver to IFM)
- **XRS_L index** 0.1-0.8 nm modeled X-ray (driver to ABBYNormal model)
- time granularity (1-minute), cadence (15-minute)




SPACE ENVIRONMENT TECHNOLOGIES

Space Weather Division

SDO-EVE workshop

SpaceWx app

<http://SpaceWx.com>



iPhone *SpaceWx* app requires solar inputs




App is an education and partnering tool:


- *SpaceWX* iPhone public education/partnership-building app (v1.2) released at Apple on August 30, 2009
- 4 upgrades covering solar, heliosphere, magnetosphere, thermosphere/ionosphere space weather domains
 - v1.3 (2009/11/12)
 - v1.4 (2010/05/01)
 - v1.5 (2010/07/06)
 - ✓ 11 partnering institutions
 - ✓ 48 datasets
 - v1.6 (2010/10/31)
 - ✓ 16 partnering institutions
 - ✓ 123 datasets
- 1345 paid downloads as of Sep 19 2010 in 39 countries

<http://SpaceWx.com>



SpaceWx output





SpaceWx Operations Recommendations

SpaceWx: solar irradiance datasets for visualization (EPO/partnering/monitoring)



- **SDO EVE** (SAM, 0.1-7 nm, 171, 257, 304, 366, Lyman-alpha, S10, XRS-B)



- **SDO AIA** (1600, 171, 193, 211, 304, 335, 94 nm and composite)



- **SET** (Mg II composite, forecasts, spectra, indices)



- **Other** (NOAA GOES, EUVS, solar cycle, proxies, SOHO, STEREO, NSO)
- time granularity (multiple), cadence (multiple)



SPACE ENVIRONMENT TECHNOLOGIES


Space Weather Division


SDO-EVE workshop

SET forecasts



SET flare probability forecast






SET GEO charging and deep dielectric discharge probabilities

GOES-10 Dec 01 2008

GOES10 135.000W geosynchronous charging environment (Sunlight)

- is negative charge
+ is positive charge
spiraling open circles:
3 hr sat intervals

Current epoch:
Dec 01 2008 18:00 UT




SDO Oct 07 2010

SDO 102W geosynchronous charging environment (shaded)

- is negative charge
+ is positive charge
spiraling open circles:
3 hr sat intervals


Current epoch:
Oct 07 2010 09:00 UT



**Charging uses 3-hour
ap in GEOPOT08**

<http://SpaceWx.com>


**Deep dielectric
discharge uses past 24
hours GOES 0.6MeV e⁻**



SET flare evolution prediction

Create X_{b10} and X_{hf}


Oct–Nov 2003 background and flares






SET flare evolution prediction

Find rate change of X-ray flux






SET flare evolution prediction

Flare rise time versus magnitude


dX_{hf}/dt vs X_{hf} for large 1 hr flares






SET flare evolution prediction

Flare magnitude & duration model





SET flare evolution prediction

Flare evolution prediction: X_{b10} , X_{hf}

