

Effective Viscosity in Plasmas

Riddhi Bandyopadhyay

Princeton University


6 April 2021

With collaborators: William H. Matthaeus, Yan Yang, Alexandros Chasapis, Vadim Roytershteyn


Background

- In collisional media (MHD), viscosity (ν) and resistivity (μ) dissipate.
- $J = \mu E$; $J \cdot E \sim J^2$
- Similarly, $\Pi_{ij} = \nu D_{ij}$; $\Pi_{ij} D_{ij} \sim D_{ij} D_{ij}$
- What about (weakly-collisional) plasmas?

J.E in Plasmas



Wan et al. (2016) PoP



Chasapis et al. (2018) ApJL

Pi-D

$$\partial_t f_\alpha + \mathbf{v} \cdot \nabla f_\alpha + \frac{\mathbf{F}}{m_\alpha} \cdot \nabla_v f_\alpha = 0 \quad \leftarrow \text{Assume Vlasov}$$

Dist. func. \downarrow $\alpha = \text{proton, electron, ...}$

$$\partial_t \mathcal{E}_\alpha^f + \nabla \cdot (\mathcal{E}_\alpha^f \mathbf{u}_\alpha + \mathbf{P}_\alpha \cdot \mathbf{u}_\alpha) = (\mathbf{P}_\alpha \cdot \nabla) \cdot \mathbf{u}_\alpha + n_\alpha q_\alpha \mathbf{E} \cdot \mathbf{u}_\alpha. \quad (1)$$

Caveat: Ignore collisions

$$\partial_t \mathcal{E}_\alpha^{th} + \nabla \cdot (\mathcal{E}_\alpha^{th} \mathbf{u}_\alpha + \mathbf{h}_\alpha) = -(\mathbf{P}_\alpha \cdot \nabla) \cdot \mathbf{u}_\alpha. \quad (2)$$


$$\partial_t \mathcal{E}^m + \frac{c}{4\pi} \nabla \cdot (\mathbf{E} \times \mathbf{B}) = -\mathbf{E} \cdot \mathbf{J} \quad (3)$$

Add internal energy: dissipation



- Integrate conservation eqns over arbitrary volume: transport on surface (or may vanish)
- Triangle diagram describes energy conversion

$$\begin{aligned} \partial_t \langle E_\alpha^f \rangle &= \langle (\mathbf{P}_\alpha \cdot \nabla) \cdot \mathbf{u}_\alpha \rangle + \langle n_\alpha q_\alpha \mathbf{E} \cdot \mathbf{u}_\alpha \rangle, \\ \partial_t \langle E_\alpha^{th} \rangle &= -\langle (\mathbf{P}_\alpha \cdot \nabla) \cdot \mathbf{u}_\alpha \rangle, \leftarrow \text{Heating!} \\ \partial_t \langle E^m \rangle &= -\langle \mathbf{E} \cdot \mathbf{j} \rangle. \end{aligned}$$

J.E and Pi-D measure somewhat different stages of energy conversion (dissipation)





Pi-D in MMS



Bandyopadhyay et al. (2021) in prep.



Pi-D in 2.5D PIC: I

Dim	L	Grid	m_i/m_e	B_0	ppg
2.5D	$102.4d_i$	8192^2	25	5.0	300



Pi-D in 2.5D PIC: II



Dim	L	Grid	m_i/m_e	B_0	ppg
2.5D	$149.6d_i$	4096^2	25	1.0	3200



Pi-D in 3D PIC

Dim	L	Grid	m_i/m_e	B_0	ppg
3D	$41.9d_i$	2048^3	50	0.5	100

Roytershteyn et al. (2015) RSPTA



Summary

- Even in plasmas $J \cdot E \sim J^2$ and $\Pi_{ij} D_{ij} \sim D_{ij} D_{ij}$
- Evaluate ν , μ for natural systems?