



# Can Coronal Dimmings be Used for Forecasting CMEs?

Khara Lukancic<sup>1,2</sup>, Rachel A. Hock<sup>1</sup>, Thomas N. Woods<sup>1</sup>, Alysha Reinard<sup>3</sup>


<sup>1</sup>Laboratory for Atmospheric and Space Physics, University of Colorado Boulder, <sup>2</sup>Metropolitan State College of Denver,

<sup>3</sup>National Oceanic and Atmospheric Administration



## Motivation:

The project goal this summer was to study the parameters of flares that exhibit coronal dimming and coronal mass ejections to better understand the relationship between coronal dimming and CME properties.




## Process:

I wrote an IDL program that analyzed 55 flares that have coronal dimming mainly looking at three specific Iron light curves using SDO-EVE data. The three Iron lines I used are Fe IX (17.1 nm, 0.65 MK), Fe XII (19.5 nm, 1.35 MK), and Fe XIV (21.1 nm, 1.86 MK). In my analysis, I found that Fe IX had the most dimming and that Fe XIV showed the least dimming. My program matches the times and locations of the flares with the CMEs as observed by SOHO and STEREO. Of the 55 flares, 34 had CMEs, and 10 had full CME catalog entries. Then, I ran correlations between the flare parameters and the CME parameters to see which had the strongest relationships.

## Flares:

When given a flare id, the code extracts the EVE data for Fe IX, Fe XII, Fe XIV, and plots the three light curves. The flare parameters are then calculated.

- ✧ **Duration** is calculated by the defined start time of the flare by GOES x-ray and the end of the flare is manually selected.
- ✧ **Depth** is calculated using the pre-flare irradiance as a baseline and finds the minimum within the flare duration.
- ✧ **Height** is calculated by using the pre-flare irradiance as a baseline and finds the maximum prior to the GOES x-ray peak.
- ✧ **Slope** is calculated by applying a linear fit to the data from the GOES x-ray peak to one-half of the dimming depth.




## CMEs:

The code finds CME matches based on location and time of the flare. The program takes the flare location and determines the expected CME location for SOHO and STEREO. Using the CME catalogs, it finds a CME within 45° of the expected location and has a CME start time within 3 hours of the flare start time. LASCO is used for limb events and STEREO A and B are used for disk-center flares.

## CME Parameters:


- ✧ Angular Width
- ✧ Velocity
- ✧ Mass (see future plans)



## Correlations and Results:

Correlations were ran between the parameters of Fe IX, Fe XII, and the CME parameters. The correlations that were discovered are justified because the slope of the dimming is related to the initial velocity of the CME. This relationship is present because the steeper the dimming slope means the dimming is occurring faster and that the material of the CME is expected to be moving faster. The depth of the dimming is expected to be related to CME mass, but this analysis is not complete (see future plans). CME mass and velocity are correlated, so the result that the depth is correlated with velocity is reasonable.

	Angular Width	Average Velocity	2nd Order Initial Velocity	2nd Order Final Velocity	2nd Order Velocity at 20R
Height Fe IX	-0.07	-0.27	-0.23	-0.21	-0.29
Depth Fe IX	0.4	0.19	0.4	<b>0.59</b>	0.08
<b>Slope Fe IX</b>	-0.02	-0.15	<b>-0.82</b>	<b>-0.56</b>	<b>-0.57</b>
<b>Duration Fe IX</b>	0.11	0.44	<b>0.54</b>	0.48	0
Height Fe XII	0.04	-0.27	-0.43	-0.47	0
Depth Fe XII	0.35	-0.08	0.01	0.44	-0.3
<b>Slope Fe XII</b>	-0.21	-0.17	<b>-0.73</b>	<b>-0.71</b>	-0.44
Duration Fe XII	0.19	-0.04	0.19	0.48	-0.36



The best found correlations are with CME second order initial velocity. The highest correlations were between Fe IX slope and Fe XII slope with CME second order velocity.

## Future Plans:

- ✧ For my senior project, I plan to focus on calculating the CME parameters of mass and velocity.
- ✧ There is high confidence in the SDO-EVE data for the Iron parameters, this is why I've chosen to focus on calculating CME parameters. There are just not enough CME data at this time to have an accurate forecast of CMEs from coronal dimming measurements. But, there is high promise to use coronal dimming as a forecast tool in the near future!

Data courtesy of NASA/SDO and the AIA & EVE teams. EVE data are available at <http://lasp.colorado.edu/eve>. AIA data are available at <http://jsoc.stanford.edu>. CME speeds courtesy of the CACTus catalog (<http://sidc.oma.be/cactus>) and the CDAW catalog ([http://cdaw.gsfc.nasa.gov/CME\\_list](http://cdaw.gsfc.nasa.gov/CME_list)).