TIMED Solar EUV Experiment: Phase E Progress Report for 2010



Submitted for NASA Grant NNX07AB68G
by Tom Woods (SEE PI)
LASP / University of Colorado
1234 Innovation Drive
Boulder, CO 80303
Phone: 303-492-4224
E-mail: tom.woods@lasp.colorado.edu
Web: http://lasp.colorado.edu/see/

SEE Science Team

LASP/CU: Tom Woods (PI), Frank Eparvier, Don Woodraska, Rachel Hock (grad. student)

NRL: Judith Lean

HAO/NCAR: Stan Solomon, Ray Roble

NASA/GSFC: Phil Chamberlin

SET: Kent Tobiska

Virginia Tech: Scott Bailey

Report Outline

- SEE Instrument Operations and Instrument Status
- SEE End of Mission Plan
- SEE Data Products
- SEE Science Overview and Recent Results
- List of Recent SEE-related Talks & Papers
- Conclusions and Future Plans for SEE

SEE Measures the Solar VUV Irradiance





EGS = EUV Grating Spectrograph

Rowland-circle grating spectrograph with 64x1024 CODACON (MCP-based) detector

XPS = XUV Photometer System

Set of 12 Si photodiodes - 8 for XUV, 1 for Ly-α, and 3 for window calibrations

XUV	EUV	FUV

EGS 27-194 nm with Δλ=0.4 nm

XPS 0.1-34 nm with Δλ=7-10 nm and Ly-α (121.6 nm) with Δλ=2 nm

FUV = Far UltraViolet: 115-200 nm
EUV = Extreme UltraViolet: 30-115 nm
XUV = X-ray UltraViolet: 0-30 nm

EGS = EUV Grating Spectrograph
XPS = XUV Photometer System

Overview of SEE Operations
and
Status of SEE Instrument

Summary of SEE Flight Operations

- Planned Experiments (through Oct 3, 2010)
  - Number of normal solar experiments = 44,981
- Actual Experiments (through Oct 3, 2010)
  - Number of normal solar experiments = 43,811 (97%)
- Calibration rockets provide degradation rates for SEE
  - NASA 36.192 launched on Feb. 8, 2002, complete success
    - Rocket results incorporated into Version 6 data
  - NASA 36.205 launched on Aug. 12, 2003, complete success
    - Rocket results incorporated into Version 7 data
  - NASA 36.217 launched on Oct. 15, 2004, complete success
    - Rocket results incorporated into Version 8 data
  - NASA 36.233 launched on Oct. 28, 2006
    - Partial success (only 0.1-36 nm and 121.6 nm irradiance measured)
  - NASA 36.240 launched on April 14, 2008, complete success
    - Rocket results incorporated into Version 10 data
  - NASA 36.258 (SDO EVE rocket) launched on May 3, 2010, complete success
    - Rocket results consistent with Version 10; no update to SEE degradation trend

List of SEE Data Gaps - Very Few Gaps

Date	State	Sensor(s)	Science Data Affected
March 1, 2002	Safe Mode	Both	Part day
March 2, 2002	Safe Mode	Both	All day
March 4, 2002	Ground SW Anomaly	EGS	All day
March 5, 2002	Ground SW Anomaly	EGS	Part day
March 19, 2002	Safe Mode	Both	Part day
March 29, 2002	Safe Mode	Both	Part day
July 24 - 30, 2002	XPS Filter Wheel Anomaly	XPS	All days
Nov. 18-19, 2002	Leonid Safing	Both	Part day
Sept. 16 - 21, 2004	TIMED Flight Software Load	Both	Sept. 16,21: Part day Sept. 17-20: All day
Sept. 29 - Oct. 1, 2004	TIMED Flight Software Load	Both	Sept. 29, Oct. 1: Part day Sept. 30: All day
May 4, 2005	Lost data due to HK rate being at 5 sec (normally 15 sec)	Both	Part day (after SSR allocation reached)

List of SEE Data Gaps - 2

Date	State	Sensor(s)	Science Data Affected
Aug. 16-18, 2006	Safe Mode	Both	Partial day on 16th All day on 17th Partial day on 18th
July 25-26, 2007	Safe Mode	Both	Partial day
Nov. 19, 2007	Safe Mode	Both	Partial day
Jan. 8-15, 2008	Safe Mode	Both	Partial day on the 8th All day from 9-15
May 24-25, 2008	Safe Mode	Both	Partial day on 24th All day on the 25th
July 7, 2008	Planning Anomaly	Both	Partial day
June 14-23	Safe Mode	Both	All days from 14-18 Partial day on the 23th
Dec. 2-3, 2009	Safe Mode	Both	All day on the 2nd Partial day on 3rd

Status of SEE Instruments

No recent changes for SEE

- **EUV Grating Spectrograph (EGS) - fully functional**
  - The EUV ($\lambda < 115$ nm) has degradation mostly at the bright lines on the CODACON (MCP-based) detector, but it is being tracked with on-board redundant channel and flat-field detector lamp weekly experiments
  - The FUV (115-195 nm) has small recovery rate that is corrected using UARS, SORCE, and XPS comparisons
- **XUV Photometer System (XPS) - 3 channels functional**
  - Fully functional until 2002/205 when there was a filter wheel anomaly (filter wheel stuck in position 6)
  - Three channels providing solar measurements
  - No spectral gaps in the XUV though because of new XPS Level 4 algorithm
- **Microprocessor Unit (MU) - fully functional**
- **SEE Solar Pointing Platform (SSPP) - fully functional**

Potential Life Issues for SEE

- EGS (grating spectrograph)
  - MCP-based detector has significant degradation at a few wavelengths (~5% of spectral range). Accuracy already degraded at those wavelengths. Degradation has slowed down with time, but still expect this degradation to continue during extended mission.
  - No degradation or anomalies for HV supply or slit changer mechanism; expect them to perform well for several more years
- XPS (set of photometers)
  - None: filter wheel mechanism is not used anymore
  - Lower priority than EGS as have SORCE XPS
- SSPP (pointing platform)
  - No degradation or anomalies for SSPP; expect it to perform well for several more years

SEE End of Mission Plan

SEE End of Mission Plan

- Overlap with SDO EVE
  - SDO EVE normal operations began May 1, 2010
    - EVE measurements are from 0.1-105 nm and at 121.6 nm
  - SDO EVE calibration rockets: May 3, 2010 and Feb. 22, 2011
  - Prefer 1-year overlap of SEE with EVE, but have limited funds left
- SEE turn-off plan
  - Continue normal SEE operations until March 1, 2011 so can overlap with SDO EVE second calibration rocket (Feb 22)
  - Perform special SEE calibrations for a month (March 2011)
    - Daily EGS calibration channel measurements, flatfield images, solar images
  - Move SSPP to 190° and turn off SEE on April 1, 2011
- SEE final data product will be version 11 (Oct. 2011 deadline)
  - Evaluate and update degradation trend over full mission
  - Evaluate and update EGS calibration in the 27-40 nm range
  - Evaluate and update XPS spectral model (7-40 nm) with EVE spectra
  - Evaluate and update EGS spectral model (114-129 nm) with SOLSTICE

SEE Data Products

SEE Version 10 Data

- Version 10 initially released in September 2009 and daily updates provided with 4-day latency
  - EGS revisions
    - Vastly improved FOV correction now uses long-term information
      - Corrects small daily jumps
    - Smaller degradation bins help with heavily degraded regions
    - Improved Gain correction now possible because of the better FOV algorithm
    - Updated FUV degradation rates from comparison to SORCE
    - Included updated EUV degradation rates using latest cal rocket (Apr 2008)
  - XPS revisions
    - Updated radiometric calibrations and updated XUV degradation rates
    - Improved empirical Gain correction
- **LASP Interactive Solar IRradiance Datacenter (LISIRD)**
  - Relatively new data center at LASP for its solar irradiance data products
    - SME, UARS SOLSTICE, TIMED SEE, SORCE, rocket experiments
    - Future missions: Glory TIM, SDO EVE
  - http://lasp.colorado.edu/LISIRD/

Summary of SEE Data Products

http://lasp.colorado.edu/see/

- Download data for individual days or merged set for the full mission
- Download IDL read / plot code
- Plot / browse data (ION script interface)

Data Product	Period	Description
SEE L2A SpWx	Orbit	8 solar indices (emissions/bands) for SpWx Ops
SEE L3	Day	1-nm spectrum from 0.5 nm to 194.5 nm, 38 emission lines, XPS 9 bands
SEE L3A	Orbit	Same as L3 but for orbit average (3-min avg)
EGS L2, L2A	D & O	0.1-nm spectrum from 27 nm to 195 nm
XPS L2, L2A	D & O	XPS 9 bands
XPS L4, L4A	D & O	0.1-nm spectral model from 0 to 40 nm
EGS L2B (Occ)	Orbit	Atmospheric transmission (single altitude)
Composite Ly-α	Day	H I Lyman-α irradiance from 1947 to present

Future SEE Data Products

- No new SEE data products are planned
- Future SEE data versions
  - Version 11 (after end of operations)
    - Expected final release will include overlap measurements with SDO-EVE and rocket underflights
    - Final clean-up of production processing code
    - Prepare products and code for delivery to TIMED archive center

SEE Science Overview and Recent Results

SEE Science Plans

Solar UV Irradiance Measurements



Obj. #1

Validations

Internal Calibrations,
Underflight Calibrations
SOHO, SNOE,
UARS, SORCE

Eparvier, Woods, Bailey, Rottman

Obj. #2

Solar UV Variability

Function of wavelength
Over time scales of minutes to years

All

Obj. #4

Modeling Solar Variation

Study variations related to active region evolution derived from solar images
Improve the **NRLEUV, SOLAR2000,** and **SunRise** solar irradiance models

Lean, Tobiska, Chamberlin, Woods

Obj. #3

Obj. #5

Study Earth's Response

Photoelectron analysis with FAST data and using the ***glow*** model
Atmospheric response studies using HAO's **TIM-GCM**

Solomon, Roble, Bailey, Eparvier

Overview of SEE Science Objectives

1. Accurately and precisely determine the time-dependent solar vacuum ultraviolet (VUV: below 200 nm) spectral irradiance.
2. Study solar VUV variability (27-day rotations, solar cycle changes) and its sources.
3. Study the solar-terrestrial relationships utilizing atmospheric models, primarily the TIME-GCM at HAO/NCAR.
4. Improve proxy models of the solar VUV irradiance
5. Determine the thermospheric neutral densities (O_2, N_2 and O) from solar occultations.

Summary of SEE Results

- **Objective 1: solar VUV spectral irradiance measurements**
  - Daily measurements since Jan. 22, 2002 with very few gaps
  - Additional analysis / validation for SEE results over full TIMED mission [Eparvier, Lean, Woods]
- **Objective 2: solar variability**
  - Updated results on solar rotation and solar cycle variations [Lean, Woods, Tobiska]
  - Updated results on flare variability as SEE for space weather research [Tobiska, Chamberlin]
    - Flare catalog on-line to make access to SEE's flare data quick and easy
- **Objective 3: model solar response in Earth's atmosphere**
  - Use of HAO TIME-GCM for atmospheric response to SEE's solar input [Solomon, Qian, Lu]
  - Use of SEE solar data and FAST photoelectron data [Peterson, Richards]
  - Comparison of GUVI FUV airglow data with SEE results [Meier, Lean, Woods]
- **Objective 4: solar irradiance modeling**
  - SOLAR2000 (S2K) model improvements [Tobiska]
  - NRLEUV model improvements [Lean, Warren]
  - Flare Irradiance Spectral Model (FISM) improvements [Chamberlin]
    - SEE data used to parameterize FISM - daily components and 1-min flare components
- **Objective 5: atmospheric density from solar occultations**
  - EGS Level 2B solar occultation data product [Eparvier]

Present, Past and Future Variations in Solar Extreme Ultraviolet Irradiance

JGR 2010 paper by J. L. Lean, T. N. Woods, F. G. Eparvier, R. R. Meier, D. J. Strickland, J. T. Correira, and J. S. Evans

SEE observations start at solar cycle maximum in 2002, include the cycle minimum in 2008-2009, and detect the rise of new cycle 24.



Paper Figure 1. Time series of daily values of a) total EUV irradiance (0-120 nm) measured by SEE are compared with b) the *Mg II index and c) the F10.7 index of solar activity for the duration of* the TIMED mission. The arrows indicate times of rocket under flight calibrations of the SEE instrument.

Present, Past and Future Variations in Solar Extreme Ultraviolet Irradiance

JGR 2010 paper by J. L. Lean, T. N. Woods, F. G. Eparvier, R. R. Meier, D. J. Strickland, J. T. Correira, and J. S. Evans

NRLEUV irradiance model compares well to SEE observations at some wavelengths.

NRLSSI 2C and 3C models developed using the SEE measurements.



Paper Figure 3. Shown are all daily solar EUV spectra in 1 nm bins measured by SEE and estimated by a) the NRLSSI 2C proxy model and b) the NRLEUV model during the TIMED mission from 2002 to 2009.

Present, Past and Future Variations in Solar Extreme Ultraviolet Irradiance

JGR 2010 paper by J. L. Lean, T. N. Woods, F. G. Eparvier, R. R. Meier, D. J. Strickland, J. T. Correira, and J. S. Evans



SEE measured solar cycle variability is larger than NRLEUV and NRLSSI 2C estimated variability at most wavelengths.

Paper Figure 5. Estimates of the solar cycle amplitudes of the EUV spectral irradiance in 1 nm bins observed by SEE and modeled by NRLSSI and NRLEUV are compared in a) as percentage increases from solar cycle minimum to maximum ((max-min)/min×100) at the two times indicated by the grey vertical lines in Figure 4. The same changes are shown in b) in energy units.

Present, Past and Future Variations in Solar Extreme Ultraviolet Irradiance

JGR 2010 paper by J. L. Lean, T. N. Woods, F. G. Eparvier, R. R. Meier, D. J. Strickland, J. T. Correira, and J. S. Evans

SEE solar cycle minimum results agree reasonably well with model estimates derived from quiet Sun radiance observations.



Paper Figure 8. The solar EUV spectral irradiance in 1 nm bins observed by SEE during solar activity minimum in August 2008 is compared in a) with the spectrum of the quiet Sun (i.e., absent magnetic activity) obtained from the NRLEUV and NRLEUV2 models. Ratios of the NRLEUV and NRLEUV2 quiet sun spectra to the SEE solar minimum spectrum are shown in b). The thick broad lines in b) are the ratios in 10 nm bins.

Present, Past and Future Variations in Solar Extreme Ultraviolet Irradiance

JGR 2010 paper by J. L. Lean, T. N. Woods, F. G. Eparvier, R. R. Meier, D. J. Strickland, J. T. Correira, and J. S. Evans



Latest versions of GUVI and SEE data agree better than ever.

Paper Figure 9. Estimates of *QEUV, the solar EUV energy < 45 nm, measured directly by SEE (one 3-*minute measurement per orbit) and derived from GUVI measurements of the FUV dayglow (orbit-averaged from sub-second measurements wherever there is dayglow along the orbit) are compared in a) as daily means for the duration of the TIMED mission. In b) and c) orbital averages are compared during 2003, when solar activity was relatively high, and in d) and e) during 2005, when solar activity levels were more moderate. The comparisons during the two 180-day periods shown in b) and d) illustrate the details of rotational modulation and the comparisons during the two 25-day periods shown in c) and e) illustrate higher resolution temporal detail and flare irradiance variations superimposed on the overall rotational modulation.

Present, Past and Future Variations in Solar Extreme Ultraviolet Irradiance

JGR 2010 paper by J. L. Lean, T. N. Woods, F. G. Eparvier, R. R. Meier, D. J. Strickland, J. T. Correira, and J. S. Evans

NRLSSI model, based on SEE observations, used to estimate past variations of the solar EUV irradiance.







Paper Figure 10. Reconstruction of daily total solar EUV irradiance (0-120 nm) with the 2C and 3C models from 1950 to 2010 are shown in a). Scenarios for annually averaged EUV irradiance changes since the Maunder Minimum derived using the 2C and 3C models are shown in b) and c). The reconstructions that incorporate background irradiance changes inferred from *Wang et al. [2005], shown by the black lines in b) and c), are the basis for the* numerical values listed in Table 1 for the Maunder Minimum.

Present, Past and Future Variations in Solar Extreme Ultraviolet Irradiance

JGR 2010 paper by J. L. Lean, T. N. Woods, F. G. Eparvier, R. R. Meier, D. J. Strickland, J. T. Correira, and J. S. Evans

Forecasting the future solar EUV irradiance level is now more accurate.



Paper Figure 12. Root-mean-square errors of the EUV irradiance changes forecast for 1 to 10 days ahead, obtained as averages of the forecast changes made over the period 2002 to 2009 using a third order autoregressive process, are compared with errors of analogous forecasts made using climatology and persistence. Approximate percentage errors of the forecasts are shown on the right ordinate.

Anomalously low solar EUV irradiance and thermospheric density during solar minimum

GRL 2010 paper by S. C. Solomon, T. N. Woods, L. V. Didkovsky, and J. T. Emmert

Solar EUV 26-34 nm irradiance is 15% ± 6% lower in 2008 than in 1996.



Paper Figure 1. Solar EUV variation in the 26–34 nm band over solar cycle 23 measured by the SEM detector on the SOHO spacecraft. Black dots: daily average values. Blue line: 81-day centered mean (with data gaps interpolated). Estimated uncertainty is 6%. Also plotted, with estimated uncertainties, are irradiance measurements integrated over the same band from the ATLAS-3 calibration rocket on 15 May 1997 (orange), the SDO/EVE EUV SpectroPhotometer (ESP) prototype (cyan) and the SDO/EVE Multiple EUV Grating Spectrograph (MEGS) prototype (magenta) on a rocket on 14 April 2008, and the TIMED/SEE 2008 annual average (green). The decrease in SEM irradiance from 1996 to 2008 is ~15%.

Anomalously low solar EUV irradiance and thermospheric density during solar minimum

GRL 2010 paper by S. C. Solomon, T. N. Woods, L. V. Didkovsky, and J. T. Emmert

Thermospheric density at 400 km is 28% lower in 2008 than in 1996. Of this, 2-5% appears to be due to higher greenhouse CO_2 concentration.



Paper Figure 3. (a) Global mean thermospheric density at 400 km altitude, obtained from satellite orbital parameters over four solar cycles. Blue: 81-day centered running mean. Black: annual average. Green dotted lines: envelope of expected decrease due to increasing CO_2 levels, in the range of 2% to 5% per decade, starting with the 1976 annual average (see text). (b) Global mean thermospheric density annual average plotted as a function of the 26–34 nm solar EUV irradiance annual average measured by the SEM for the ascending (red) and descending (blue) phases of solar cycle 23.

Anomalously low solar EUV irradiance and thermospheric density during solar minimum

GRL 2010 paper by S. C. Solomon, T. N. Woods, L. V. Didkovsky, and J. T. Emmert



TIM-GCM model results confirm that lower thermospheric density is consistent with lower solar EUV irradiance in 2008-2009.

Paper Figure 4. Thermospheric temperature and density modeled by the NCAR TIE-GCM on day-of-year 227 using the spectra shown in Figure 2. (a, b) Model temperature and density at 400 km for 1996. (c, d) Model temperature and density for 2008. (e, f) Global average temperature and density as a function of altitude for 1996 and 2008. Black line: 1996. Red line: 2008 with both decreased solar EUV and increased CO_2. (g, h) Global average temperature change from 1996 to 2008, and density ratio for 2008 divided by 1996, as a function of altitude. Black line: 1996. Blue line: 2008 with only solar EUV decrease. Red line: 2008 with both decreased solar EUV and increased CO_2.

Recent SEE-related Talks & Papers

Recent SEE Related Talks

- AGU, December 2009, 3 talks
- Boulder Solar Day, March 2010, 1 talk
- AAS / SPD, May 2010, 2 talks
- Aspen Global Change Institute (AGCI Workshop), June 2010, 1 talk
- COSPAR, July 2010, 4 talks

Recent SEE Related Papers

- Solomon, Stanley, T. Woods, L. Didkovsky, J. T. Emmert, and L. Qian, Anomalously low solar extreme-ultraviolet irradiance and thermospheric density during solar minimum, *Geophysical Research Letters,* 37, 16, 2010.
- Tobiska, W. K. and S. D. Bouwer, Distributed networks enable advances in U.S. space weather operations, *J. Adv. Space Res*., 10.1016/j.asr.2010.07.009, 2010.
- Tobiska, W. K, G. Crowley, S. J. Oh, and M. Guhathakurta, Space Weather Gets Real—on Smart Phones, Space Weather, in press paper #2010SW000619, 2010.
- Lean, J. L., T. N. Woods, F. Eparvier, R. R. Meier and D. J. Strickland, Solar EUV Irradiance, Past, Present and Future, *J. Geophys. Res.,* in press, 2010.
- Lu, G., M. G. Mlynczak, T. Woods, and R. G. Roble, On the relationship of Joule heating and nitric oxide radiative cooling in the thermosphere, *J. Geophys. Res.,* 115, A05306, 2010.
- Del Zanna, G., V. Andretta, P. C. Chamberlin, T. N. Woods, and W. T. Thompson, The EUV spectrum of the Sun: SOHO CDS NIS radiometeric calibration, *Astron. & Astrophys*., in press, 2010.
- Lean, Judith L., and Thomas N. Woods, Solar Spectral Irradiance Measurements and Models, in Evolving Solar Physics and the Climates of Earth and Space, Karel Schrijver and George Siscoe (Eds), Cambridge Univ. Press, 2010.

Conclusions and Future Plans

Summary of SEE Observations

- TIMED SEE has been very successful in obtaining new, accurate measurements of the solar EUV irradiance
  - SEE data available from http: //lasp.colorado.edu/see/
- More than 100 flares have been observed by SEE
  - Extreme flare periods are April 2002, July 2002, May-June 2003, Oct.-Nov. 2003, July 2004, Jan. 2005, Sept. 2005, and Dec. 2006
  - Large flares vary as much as 11-year solar cycle variations
  - New flare models have been developed with SEE observations
- More than 120 solar rotations have been observed by SEE
  - Variability of 5-70% observed (wavelength dependent)
- TIMED mission has observed solar maximum and minimum activity during solar cycle 23 and the start of cycle 24 in 2009

SEE Plans for 2011

- Plan to terminate the SEE operations in April 2011
  - Deactivate SEE after calibration rocket flight planned for Feb. 22, 2011
  - See slide 11 for more details
- Produce final data product for SEE by October 2011
  - Final SEE product will be version 11
  - See slide 11 for more details